

9-10-2013

Anti-chaperone Activity and Cytotoxicity of Chemical Components in Copaiba Oil

Daniel Kulman
Cleveland State University

Janine Naim
Cleveland State University

Bin Su Ph.D.
Cleveland State University, B.SU@csuohio.edu

Follow this and additional works at: https://engagedscholarship.csuohio.edu/u_poster_2013

 Part of the [Macromolecular Substances Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Kulman, Daniel; Naim, Janine; and Su, Bin Ph.D., "Anti-chaperone Activity and Cytotoxicity of Chemical Components in Copaiba Oil" (2013). *Undergraduate Research Posters 2013*. 23.
https://engagedscholarship.csuohio.edu/u_poster_2013/23

This Article is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2013 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

Anti-chaperone activity and cytotoxicity of chemical components in Copaiba oil

College of Sciences and Health Professions

Student Researchers: Daniel Kulman; Janine Naim

Faculty Advisor: Bin Su, Ph.D.

Abstract

Copaiba oil derived from the oleoresin of the Copaiba tree has been widely used as an antiseptic and expectorant for the respiratory tract, and as anti-inflammatory agent in various skin diseases. Studies have indicated that Copaiba oil exhibited anti-carcinogenic properties in various preclinical studies. However, the anti-cancer mechanisms of copaiba oil still remain unclear. There are various diterpenoid compounds within Copaiba oil, which also make the mechanism investigation very difficult. Hardwickiic acid (HAA), a clerodane diterpenoid isolated from Copaiba oil shows anti-chaperone activity from a recent study. In the current study, cytotoxicity and anti-chaperone assay guided isolation led to 9 fractions from Copaiba oil. Three of the fractions showed cytotoxicity in prostate cancer cells. And other three fractions exhibited potent anti-chaperone activity. There are multiple chemical components in the fractions that showed cytotoxicity, which has been confirmed with mass spectrum. The three fractions showed anti-chaperone activity were further purified for structure elucidation. NMR combined with MS reveal that the three fractions are Copaibic acid, Hardwickiic acid and 7-Acetyl-copaibic acid. All three compounds can be used as lead compounds for the development of more potent small molecule chaperone inhibitors.